

**For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory**

### **Abridged Prescribing Information**

### **DEPAKOTE® XR**

### **Divalproex Extended-Release Tablets IP**

#### **COMPOSITION**

DEPAKOTE® XR 250mg: Divalproex Sodium I.P. equivalent to Valproic Acid 250 mg

DEPAKOTE® XR 500mg: Divalproex Sodium I.P. equivalent to Valproic Acid 500 mg

#### **THERAPEUTIC INDICATIONS**

Treatment of acute manic or mixed episodes associated with bipolar disorders with or without psychotic features. Indicated as monotherapy and adjunctive therapy in treatment of adult patients and pediatric patients (age of 10 years or older) with complex partial seizures that occur in isolation or in association with other seizure types. As sole and adjunctive therapy of simple and complex absence seizures in adults and children 10 years of age or older and adjunctively in adults and children 10 years of age or older with multiple seizure types that include absence seizures.

**DOSAGE AND ADMINISTRATION :** Mania : Recommended initial dose is 25mg/kg/day, once daily. Maximum recommended dosage is 60mg/kg/day. Epilepsy : Complex Partial Seizures (For adults and children 10 years of age or older) : Monotherapy (Initial Therapy) with 10-15mg/kg/day. Conversion to Monotherapy : Initiate therapy at 10-15mg/kg/day. Concomitant anti-epileptic drug to be reduced by approximately 25% every 2 weeks. Adjunctive Therapy: initiate therapy at 10-15mg/kg/day. Simple and Complex Absence Seizures : Recommended initial dose is 15mg/kg/day. Maximum recommended dosage is 60mg/kg/day. Conversion from Divalproex Sodium Delayed Release to Depakote XR : Depakote XR should be administered once daily using a dose 8-20% higher than the total daily dose of Divalproex Sodium Delayed Release. Valproate treatment must be started and supervised by a doctor experienced in managing epilepsy or bipolar disorder.

**SAFETY-RELATED INFORMATION: Contraindications:** Hypersensitivity to sodium valproate or valproic acid; Acute and chronic hepatitis; Patient or family history of severe hepatitis, especially drug related; Hepatic porphyria; Patients known to have mitochondrial disorders caused by mutations in the nuclear gene encoding mitochondrial enzyme polymerase  $\gamma$  and in children under two years of age who are suspected of having a POLG-related disorder; Patients with known urea cycle disorders.

**Warnings:** Severe liver damage resulting sometimes in fatalities has exceptionally been reported, severe pancreatitis which may result in fatalities has been very rarely reported.

***Female children, female adolescents, women of childbearing potential and pregnant women:***

Valproate should not be used in female children, in female adolescents, in women of child-bearing potential and pregnant women unless alternative treatments are ineffective or not tolerated because of its high teratogenic potential and risk of developmental disorders in infants exposed in utero to valproate. Children exposed in-utero to valproate are at a high risk of serious developmental disorders (in up to 30-40% of cases) and / or congenital malformations (in approximately 10% of cases). Carefully balance the benefits of valproate treatment against the risks when prescribing valproate for the first time, at routine treatment reviews, when a female child reaches puberty and when a woman plans a pregnancy or becomes pregnant. Women of child bearing potential must use effective contraception during treatment and be informed of the risks associated with the use of valproate during pregnancy.

The prescriber must ensure that the patient is provided with comprehensive information on the risks alongside relevant materials to support her understanding of the risks. In particular the prescriber must ensure the patient understands :

- The nature and the magnitude of the risks of exposure during pregnancy, in particular the teratogenic risks and risks of developmental disorders.
- need to use effective contraception;
- need for regular review of treatment;

- the need to rapidly consult her physician if she is thinking of becoming pregnant or there is a possibility of pregnancy.

Valproate therapy must be started and supervised by a doctor experienced in managing epilepsy or bipolar disorder.

***Suicidal ideation and behavior*** : Patients should be monitored for signs of suicidal ideation and behavior and appropriate treatment should be considered.

***Carbapenem agents*** : Concomitant use of valproate and carbapenem agents is not recommended.

***Patients with known or suspected mitochondrial disease*** : Valproate may trigger or worsen clinical signs of underlying mitochondrial diseases caused by mutations or mitochondrial DNA as well as the nuclear-encoded POLG gene.

***Aggravated convulsions***

As with other antiepileptic drugs, some patients may experience, instead of an improvement, a reversible worsening of convulsion frequency and severity (including status epilepticus), or the onset of new types of convulsions with valproate. In case of aggravated convulsions, the patients should be advised to consult their physician immediately. (see *Adverse Reaction*)

**Precautions:** Liver function tests should be carried out before therapy, and periodically during the first 6 months especially in patients at risk. Blood tests are recommended prior to initiation of therapy or before surgery, and in case of spontaneous bruising or bleeding. Potential benefit should be weighed against its potential risk in patients with systemic lupus erythematosus. When a urea cycle enzymatic deficiency is suspected, metabolic investigations should be performed prior to treatment because of the risk of hyperammonemia with valproate. Patients should be warned of the risk of weight gain at initiation and appropriate strategies should be adopted to minimize the risk. Patients with an underlying carnitine palmitoyltransferase (CPT) type II deficiency should be warned of the greater risk of rhabdomyolysis when taking valproate. Alcohol intake is not recommended during treatment. Monotherapy is recommended in children under the age of 3 years. Concomitant use of salicylates should be avoided in children under 3 due to the risk of liver toxicity. Patients with renal insufficiency should be closely monitored, it may be necessary to decrease the dosage.

**Pregnancy** : Not to be used during pregnancy and in women of child-bearing potential unless clearly necessary. Women of childbearing potential should be informed of the risks and benefits of use of valproate in pregnancy.

**Lactation** : Excretion of valproate in breast milk is low with a concentration of 1% to 10% of maternal serum levels. Based on literature breastfeeding can be envisaged taking into account the valproate safety profile especially hematological disorders.

**Adverse Reactions:** Very common ( $\geq 1\%$ ): tremor, nausea, Common ( $\geq 1\%$  and  $< 10\%$ ) : anemia, thrombocytopenia, extrapyramidal disorder, stupor, somnolence, convulsion, memory impairment, headache, nystagmus, dizziness, vomiting, gingival disorder (mainly gingival hyperplasia), stomatitis, abdominal pain (upper), diarrhea, hypersensitivity, transient and /or dose related alopecia, nail and nail bed disorders, hyponatremia, weight increased, hemorrhage, liver injury, dysmenorrhea, confusional state, hallucinations, aggression, agitation, disturbance in attention.

*For full prescribing information, please contact Sanofi-Synthelabo (I) Pvt Ltd, Sanofi House, CT Survey No 117-B, L& T Business Park, Saki Vihar Road, Powai, Mumbai-400072*

Source: CCDS version 21 dated 25<sup>th</sup> August 2016, REX PI dated Sept 2011, content of the DHPC

Updated: October 2016