For the use only of a Registered Medical Practitioner or a Hospital or a Laboratory | This package insert is continually updated. Please read carefully before using a new pack

Fluticasone Furoate and Azelastine Hydrochloride Nasal Spray

Allegra®nasal Duo

Each actuation delivers: Fluticasone Furoate Azelastine Hydrochloride I.P. COMPOSITION:

Excipients

DOSAGE FORM

Allegra® nasal Duo is indicated for the treatment of symptoms of

allergic rhmitis. Allegar's massi Duo is indicated for the management of symptoms of allergic rhmitis, once the need for an antihistamine and conticosteroid has been established. It is recommended for the teathern of pensistent, moderate to severe symptoms in adults and adolescents above 12 years of age. DOSAGE AND ADMINISTRATION.

DOSAGE AND ADMINISTRATION.
Allogard insall but obtacle to administered by the intranssal rode only,
Adults and Adolescents (12 years of age and older):
One somy no roads those daily.
The recommended dosage should not be exceeded.
Shale the bottle well before each use.
For intranssal use only, Do not spray in the mouth and eyes.
CONTRAINDLATIONS.
CONTRA

This formulation is not recommended for children under 12 years.

Statemic Particianos of Tital.

Tita

caterine furnale heats scray, and architectures and topical confoosteroid all distultance may be reported with symptoms coult as fluenced vision or other and the scray of the scray of the scray of the scray of the the scray of the scray of the scray of the scray of the the scray of scray

Adulation kydrochlorios <u>CRS Decreasars</u>

<u>Concurrent use of azelasine hydrochloride</u> with alcohol or other CNS <u>CRS Decreasars</u>

<u>Concurrent use of azelasine hydrochloride</u> with alcohol or other CNS <u>Associated or CRS Decreasars</u>

<u>CRS Dec</u>

<u>Presensor</u>
<u>Fluticasone furcate</u>: There are no adequate data from the use of futiciasone furcate in pregnant women. Fluticasone furcate should be used in pregnancy only if the benefits to the mother outweigh the potential risks to the feetus or child.

the foetus or child.

Azelastine Hydrochloride: Limited data from post marketing experience over decades of use with azelastine hydrochloride in pregnant women have not identified any drug associated risks of miscarriage, birth defects, or other adverse maternal or foetal outcomes.

White using Afergain hasal Duo hasal spery driving, operating machinely, or any other fuzzionis sorthees should be induced.

ADVERSE REACTIONS

ADVERSE REACTIONS

ADVERSE REACTIONS

Linear to the classification of respective to the control of the classification of a reflect of the classification of a reflect of the classification of a reflect of the classification of the cl

Not known: Bronchospasm Musculoskeletal and connective tissue disorder (children)

Growth retardation (known)
Immune system disorders:
Rare: Hypersensitivity reactions including anaphylaxis, angioedema, rash, and urticaria . Nervous system disorders: Common: Headache

ning/burning, tearing/watering and redness of the eyes) in the studies icacy was maintained over the full 24-hour dosing period with once-daily

Conset of therapeutic benefit was observed as early as 8 hours after initial administration, with further improvement observed for several dates.

consisting proceedings of the contraction of the co

special bornay and a second se

hepatic systemane P450 isosyme, CYPSA4, the pharmacoinests of fulficacions furcina may be altered in placeins with repatic impairment. The systemic exposure would be expected to be higher than that observed had the stilly been controlled after mighty because another in patients with severe languages and the still severe higher than the controlled spray with cause in patients with severe higher impairment.

Platitistics for the still severe higher impairment.

Platitistics for the still severe higher impairment.

Platitistics for the still severe higher impairment.

Alternatives for the still severe higher impairment.

Alternative of the still severe higher impairment.

Alternative of the still severe higher impairment is a still severe higher impairment.

Alternative Sylptics for the required in publish with renal impairment.

Alternative Sylptics for the severe higher impairment.

unhanged acelestra.
Baseal Palestand acelestra.
Baseal Palestand Service of the Commission of the Co

Manufactured in field by Mic Biotel Physics Saini Majra, Nalagarh Repair Road, Nalagarh Denict Solan-174101 (H.P.), NIDA. South Majra, Nalagarh, Denict Solan-174101 (H.P.), NIDA. South Federal Majra, South Federal Chem. South Federal Chem. South Federal Chem. South Federal Chem. Park, Sain Yhar Road, Powel, Muribal –400072 Updated the 2023 Decided Chem. South Federal Chem. South Fede

Steps to use your nasal spray correctly

Parts of the Nasal Spray







then rentove the product cap.
Hold the bottle as shown with your forefinger and middle finger on either side of the nozzle and your thumb underneath the bottle.

STEP 2 Shake the bottle gently and

STEP 3
If using for the first time or if you have not used it for a week or more, test the spray. Testing the spray. With the nozzle pointing away from you, press down a few times as shown until a fine mist comes out of the nozzle.



STEP 4
Close one nostril and hold the bottle as snown in step 2.
Titl your head forward slight ly and keeping the bottle upright, carefully insert the tip of the nozzle in the other



Start to breathe in through your nose and while breathing in, press down with your fingers once to release a spray. 330mm

Breathe out through your mouth. Repeat steps 5 and 6 to inhale a second spray



STEP 7 Repeat steps 4, 5 and 6 for the other nostril.

Wipe the nozzle with a clean handkerchief / tissue and replace the protective dust cap.

IMPORTANT POINTS TO NOTE WHILE USING THE NASAL SPRAY







. For effective use of the nasal spray PRESS DOWN THE FINGER
FLANGE DISC COMPLETELY. incomplete or partial press will not release the medicine







